Author Index

Aamodt, S., see Alfredson, T. 213
Alfredson, T.
——, Sheehan, T., Lenert, T., Aamodt, S. and Correia, L.
Determination of compound identity by group type and class using diode-array detection in high-performance liquid chromatography 213
Alton, K. B.
——, Petruzzi, R. F. and Patrick, J. E.
High-performance liquid chromatographic assay for azatadine in human urine 249
Angus, L., Stein, T.A. 377
Arvidsson, T., see Daoud, N. 311
Auguste, L. J., see Stein, T.A. 377
Balistrera, W. F., see Setchell, K. D. R. 267
Bao, Z., see Mushtaq, M. 293
Benedict, C. R.
Simultaneous measurement of urinary and plasma norepinephrine, epinephrine, dopamine, dehydroxyphenylalanine, and dihydroxyphenylacetic acid by coupled-column high-performance liquid chromatography on C_8 and C_{18} stationary phases 369
Berry, D. J.
Reversed-phase high-performance liquid chromatographic determination of chloramphenicol in small biological samples 337
Biaggi, M. E., see Binder, S. R. 241
Billiet, H. A. H., see Strasters, J. K. 181
Binder, S. R.
—— and Biaggi, M. E.
Analysis of urinary catecholamines by high-performance liquid chromatography in the presence of labetalol metabolites 241
Borra, C.
——, Han, S. M. and Novotny, M.
Quantitative analytical aspects of reversed-phase liquid chromatography with slurry-packed capillary columns 75
Borrero, E., see Stein, T. A. 377
Brendel, E.
——, Meineke, I., Witsch, D. and Zschunke, M.
Simultaneous determination of 5-aminosalicylic acid and 5-acetylaminosalicylic acid by high-performance liquid chromatography 299
Brooks, H. B.
——, Thrall, C. and Tehrani, J.
High-performance liquid chromatography system for packed capillary columns 55
Brown, H., see Holland, O. B. 393
Brown, P. R., see Simpson, R. C. 41
Correia, L., see Alfredson, T. 213
Daoud, N.
——, Arvidsson, T. and Wahlund, K.-G.
Pre-column-venting plug technique for direct injection of untreated blood plasma samples into reversed-phase liquid chromatography systems 311
De Galan, L., see Strasters, J. K. 181
Dubay, G., see Hook, D. J. 99
Dube, P., see Holland, O. B. 393
Duff, R.
—— and Murrill, E.
Determination of L-buthionine-(*S*,*R*)-sulfoximine in plasma by high-performance liquid chromatography with *o*-phthalaldehyde derivatization and fluorometric detection 275
Dutta, P. K.
—— and O'Donovan, G. A.
Separation and quantitation of bacterial ribonucleoside triphosphates extracted with trifluoroacetic acid, by anion-exchange high-performance liquid chromatography 119
Fekete, J., see Horvai, G. 25
Galan, L. de, see Strasters, J. K. 181
Han, S. M., see Borra, C. 75
Hartman, P. A.
—— and Stodola, J. D.
Gradient reversed-phase high-performance liquid chromatographic separation of paldimycin (U-70,138) antibiotics and related compounds 363
Hjertén, S., see Yao, K. 87
Holland, O. B.
——, Risk, M., Brown, H., Komes, K., Dube, P. and Swann, C.
High-performance liquid chromatography in the radioimmunoassay of urinary tetrahydroaldosterone 393
Hook, D. J.
——, More, C. F., Yacobucci, J. J., Dubay, G. and O'Connor, S.
Integrated biological–physicochemical system for the identification of antitumor compounds in fermentation broths 99
Horvai, G.
——, Fekete, J., Niegreisz, Zs., Tóth, K. and Pungor, E.
Liquid chromatography with electrochemical detection. New applications in ion chromatography and reversed-phase chromatography 25

Hung, C. T., see Taylor, R. B. 383
Johansson, M.
—— and Westerlund, D.
Determination of N-acetylcysteine, intact and oxidized, in plasma by column liquid chromatography and post-column derivatization 343
Jorde, C., see Trathnigg, B. 17
Joyce III, T. H., see Wong, J. K. 261
Juergens, U.
Simultaneous determination of Zonisamide and nine other anti-epileptic drugs and metabolites in serum. A comparison of microbore and conventional high-performance liquid chromatography 233
Kabra, P. M.
—— and Wall, J. H.
Liquid chromatographic determination of cyclosporine in whole blood with the advanced automated sample processing unit 305
Kannan, R.
——, Miller, S., Verez, P. and Singh, B. N.
Sensitive method for the measurement of amiodarone and desethylamiodarone in serum and tissue and its application to disposition studies 225
Kateman, G., see Strasters, J. K. 181
Kendle, K. E., see Taylor, R. B. 383
Kirkland, J. J., see Köhler, J. 125
Klíma, J.
——, Petrásek, R. and Kočandrle, V.
Simple and specific isocratic liquid chromatographic procedure for cyclosporine A determination in whole blood, compared with radioimmunoassay 357
Klooster, M. J., see Setchell, K. D. R. 267
Kočandrle, V., see Klíma, J. 357
Köhler, J.
—— and Kirkland, J. J.
Improved silica-based column packings for high-performance liquid chromatography 125
Komes, K., see Holland, O. B. 393
Kowalski, B. R., see Ramos, L. C. 165
——, see Sanchez, E. 151
Lammert, S. A., see McFadden, W. H. 201
Lang, J. K.
—— and Packer, L.
Quantitative determination of vitamin E and oxidized and reduced coenzyme Q by high-performance liquid chromatography with in-line ultraviolet and electrochemical detection 109
Lenert, T., see Alfredson, T. 213
LeRoy, A. F., see Parsons, P. J. 323
Lim, C. K., see Setchell, K. D. R. 267
McFadden, W. H.
—— and Lammert, S. A.
Techniques for increased use of thermospray liquid chromatography–mass spectrometry 201
Meineke, I., see Brendel, E. 299
Miller, S., see Kannan, R. 225
More, C. F., see Hook, D. J. 99
Morrison, P. F., see Parsons, P. J. 323
Morrow, D. H., see Wong, J. K. 261
Murrill, E., see Duff, R. 275
Mushtaq, M.
——, Bao, Z. and Yang, S. K.
Reversed-phase high-performance liquid chromatographic separation of phenolic derivatives of benzo[*a*]pyrene, benz[*a*]anthracene, and chrysene with monomeric and polymeric C_{18} columns 293
Niegreisz, Zs., see Horvai, G. 25
Niessen, W. M. A.
—— and Poppe, H.
Open-tubular liquid chromatography–mass spectrometry with a capillary-inlet interface 1
Noguchi, K., see Seki, T. 283, 287
Novotny, M., see Borra, C. 75
O'Connor, S., see Hook, D. J. 99
O'Donovan, G.A., see Dutta, P. K. 119
Packer, L., see Lang, J. K. 109
Parsons, P. J.
——, Morrison, P. F. and LeRoy, A. F.
Determination of platinum-containing drugs in human plasma by liquid chromatography with reductive electrochemical detection 323
Patrick, J. E., see Alton, K. B. 249
Perez, V., see Kannan, R. 225
Petrásek, R., see Klíma, J. 357
Petruzzi, R. F., see Alton, K. B. 249
Poppe, H., see Niessen, W. M. A. 1
Pungor, E., see Horvai, G. 25
Ramos, L. C.
——, Sanchez, E. and Kowalski, B. R.
Generalized rank annihilation method. II. Analysis of bimodal chromatographic data 165
——, see Sanchez, E. 151
Reid, R. G., see Taylor, R. B. 383
Risk, M., see Holland, O. B. 393
Sanchez, E.
——, Ramos, L. C. and Kowalski, B. R.
Generalized rank annihilation method. I. Application to liquid chromatography–diode array ultraviolet detection data 151
——, see Ramos, L. C. 165

Schlabach, T.D.
—— and Wilson, K. D.
Microbore flow-rates and protein chromatography 65
Seki, T.
——, Noguchi, K. and Yanagihara, Y.
Determination of riboflavine in human urine by the use of a hydrophillic gel column 283
——, Yamaguchi, Y., Noguchi, K. and Yanagihara, Y.
Determination of ascorbic acid in human urine by high-performance liquid chromatography coupled with fluorimetry after post-column derivatization with benzamidine 287
Setchell, K. D. R.
——, Welsh, M. B., Klooster, M. J., Balistreri, W. F. and Lim, C. K.
Rapid high-performance liquid chromatography assay for salivary and serum caffeine following an oral load. An indicator of liver function 267
Sheehan, T., see Alfredson, T. 213
Simpson, R. C.
—— and Brown, P. R.
Development of a microbore high-performance liquid chromatographic system for biological applications 41
Singh, B. N., see Kannan, R. 225
Stein, T. A.
——, Angus, L., Borrero, E., Auguste, L. J. and Wise, L.
Program measurement of prostaglandin E_2 synthesis by gastric mucosa by high-performance liquid chromatography 377
Stodola, J. D., see Hartman, P. A. 363
Strasters, J. K.
——, Billiet, H. A. H., De Galan, L., Vandeginste, B. G. M. and Kateman, G.
Evaluation of peak-recognition techniques in liquid chromatography with photodiode array detection 181
Swann, C., see Holland, O. B. 393
Taylor, R. B.
——, Kendle, K. E., Reid, R. G. and Hung, C. T.
Chromatography of progesterone and its major metabolites in rat plasma using microbore high-performance liquid chromatography columns with conventional injection and detection systems 383
Tehrani, J., see Brooks, H. B. 55
Thrall, C., see Brooks, H. B. 55
Tóth, K., see Horvai, G. 25
Trathnigg, B.
—— and Jorde, C.
New universal detector for high-performance liquid chromatography: the density detector 17
Vandeginste, B. G. M., see Strasters, J.K. 181
Vargo, J. D.
Comparing borosilicate and soda-lime glass for fabricating open-tubular liquid chromatographic columns 33
Wahlund, K.-G., see Daoud, N. 311
Wall, J. H., see Kabra, P. M. 305
Welsh, M. B., see Setchell, K. D. R. 267
Westerlund, D., see Johansson, M. 343
Wilson, K. D., see Schlabach, T. D. 65
Wise, L., see Stein, T. A. 377
Witsch, D., see Brendel, E. 299
Wong, J. K.
——, Joyce III, T. H. and Morrow, D. H.
Determination of hydralazine in human plasma by high-performance liquid chromatography with electrochemical detection 261
Yacobucci, J. J., see Hook, D. J. 99
Yamaguchi, Y., see Seki, T. 287
Yanagihara, Y., see Seki, T. 283, 287
Yang, S. K., see Mushtaq, M. 293
Yao, K.
—— and Hjertén, S.
Gradient and isocratic high-performance liquid chromatography of proteins on a new agarose-based anion exchanger 87
Zschunke, M., see Brendel, E. 299